



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 1 of 14

Table of Contents

1. Overview 3
 Purpose.....3

2. Mice 3
 2.1 Anesthetics3
 2.2 Anesthetic Reversal Agents4
 2.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)4
 2.4 Analgesics- Opioids (controlled substances)5
 2.5 Analgesics- Local Anesthetic/Analgesics5

3. Rats 6
 3.1 Anesthetics6
 3.2 Anesthetic Reversal Agents7
 3.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)7
 3.4 Analgesics- Opioids (controlled substances)8
 3.5 Analgesics- Local Anesthetic/Analgesics8

4. Rabbits 9
 4.1 Anesthetics9
 4.2 Anesthetic Reversal Agents9
 4.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)10
 4.4 Analgesics- Opioids (controlled substances)10
 4.5 Analgesics- Opioids (controlled substances)11
 4.6 Analgesics- Local Anesthetic/Analgesics11

5. Swine..... 12
 5.1 Anesthetics12
 5.2 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)12
 5.3 Analgesics- Opioids (controlled substances)13

6. Fish..... 13
 6.1 Anesthetics13

7. Frogs..... 13



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 2 of 14

7.1 Anesthetics13

8. Resources 14

9. Revisions..... 14



Drug Formulary of Anesthetic and Analgesic Agents for Laboratory Animals

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 3 of 14

1. Overview

Purpose

Refer to ACU-211 Non-Pharmaceutical Grade Agents or Mixtures of Pharmaceutical (Cocktails) Use Procedures

<http://drexel.edu/research/animal-research/IACUC/Policies%20and%20Procedures/>

All controlled substances must be logged per dose in a drug logbook.

2. Mice

2.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Isoflurane	4-5% (induction) 1-3% (maintenance)	Inhalant	Continuous	Due to rapid induction and recovery from isoflurane, investigators must provide supplemental analgesic for invasive survival procedures prior to beginning surgical procedure; must use precision vaporizer.
Ketamine/Xylazine	80-100 (ketamine) + 5-20 (xylazine) [in same syringe]	IP	Once at induction, with additional doses as needed	Ketamine is a controlled substance . Xylazine is a controlled substance in Pennsylvania. Additional doses must be given at ½ strength. Dilute with sterile saline to make each dose volume 1.0ml.
Ketamine/Xylazine/Acepromazine	70-100 (ketamine) +10-20 (xylazine) +2-3 (acepromazine) [in the same syringe]	IP	Once at induction, with additional doses as needed	Ketamine is a controlled substance . Xylazine is a controlled substance in Pennsylvania. Additional doses must be given at ½ strength. Dilute with sterile saline to make each dose volume 1.0ml.
Sodium Pentobarbital (Nembutal)	40-50	IP	Once at induction	Controlled substance recommended for terminal/acute procedures; must provide supplemental analgesic for invasive survival procedures. Difficult to maintain. Narrow margin of safety. Not recommended for survival surgery.



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 4 of 14

Tribromoethanol (Avertin) (2.5% Solution)	250-500	IP	Once at induction	Not recommended. Nonpharmaceutical grade drug, its use must be justified in the protocol due to scientific reasons. Diluted solution must be properly stored and used within 30 days of initial preparation. *see note below
--	---------	----	-------------------	---

***AVERTIN**

100% stock Avertin (Tribromoethanol)

Mix: add tribromoethanol to tertiary amyl alcohol and dissolve by heating and stirring. Add distilled water and continue until the solution is well mixed. Store wrapped in foil (light sensitive solution, okay to use brown glass bottle), 4°C. Solution may have to be warmed to dissolve. Mixture should be clear. 10g tribromoethyl alcohol (2, 2, 2 tribromoethanol), Aldrich T4, 840-2 10ml tertiary amyl alcohol (2 methyl-2-butanol), Aldrich 24, 048-6

Warning! Decomposition can result from improper storage. 2.5% diluted Avertin solution must be used within 30 days of initial preparation and be properly stored. Be sure to label the container with the date of preparation.

For use in mice, dilute the 100% to 2.5% (1:40) using diluent, water or isotonic saline. Diluent recipe: 0.8% NaCl 1mM Tris (pH 7.4) 0.25mM EDTA Check the pH. Adjust to pH 7.4. To make 50 ml 2.5% Avertin, add 1.25 ml 100% to 48.75 ml liquid (diluent, water or saline) Filter .22 micron Store at 4° C, away from light in foil wrap or brown bottle Dosage for mice may vary with different preparations of Avertin. Dosage should be recalculated each time a 100% stock is made up. Test for best effect in a few mice before choosing dose. Allow 5-10 min to take effect.

2.2 Anesthetic Reversal Agents

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Yohimbine	1.0-2.0	SC or IP	When needing to reverse the effects of Xylazine	Yohimbine DOES NOT reverse the effects of Ketamine

2.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)

*Note: Prolonged use of NSAIDs may cause renal, gastrointestinal, or other problems



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 5 of 14

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Meloxicam (Metacam)	5-10	PO or SC	Once pre-operatively then every 8-12 hours as needed	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.
Carprofen (Rimadyl)	5-20	SC	Once pre-operatively 5mg every 12 hours, 20 mg every 24 hours	Depending on the procedure, may be used as sole analgesic, or as a multi-modal analgesia with buprenorphine.
Ketoprofen (Ketofen)	2-5	SC	Once pre-operatively then every 12-24 hours as needed	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.

2.4 Analgesics- Opioids (controlled substances)

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Buprenorphine	0.1-0.5	SC or IP	Once pre-operatively for preemptive analgesia and postoperatively every 4-6 hrs	When used as sole analgesic, typical regimen is: once at time of procedure, second dose will be administered 4-6 hours later. Additional doses every 4-6 hours as needed. Consider multi-modal analgesia with NSAID and local analgesic. Controlled substance
Buprenorphine Sustained-Release Lab (Bup SR Lab) 1mg/ml	0.5-1.0 (recommend 0.6 to start)	SC	Once pre-operatively for preemptive analgesia	Provides 48 hours of pain relief; sedation may be seen for duration of effect. DO NOT DILUTE. Controlled substance

2.5 Analgesics- Local Anesthetic/Analgesics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Bupivacaine (Marcaine, Sensorcaine)	Dilute to 0.25%; do not	SC or intradermal	Locally before making surgical incision, or before final skin closure	4-8 hours duration of action. Can be used with inhalants if other analgesics cannot be administered before animals awakened from isoflurane.



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 6 of 14

	exceed 8mg/kg total dose	surrounding incision		
Lidocaine Hydrochloride	Dilute to 0.5%; do not exceed 7mg/kg total dose	SC or intradermal surrounding incision	Locally before making surgical incision, or before final skin closure	Faster onset than Bupivacaine, but short (<1 hour) duration of action. May be combined with Bupivacaine.

Refer to ULAR Recommendations on Use of Bupivacaine for Perioperative Analgesia
[Bupivacaine \(Marcaine\) Recommendations for Use in Perioperative Analgesia](#)

3. Rats

3.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Isoflurane	4-5% (induction) 1-3% (maintenance)	Inhalant	Continuous	Rapid induction and recovery. Must provide supplemental analgesic for invasive survival procedures prior to beginning surgical procedure; must use precision vaporizer
Ketamine/Xylazine	80-100 (ketamine) + 5-10 (xylazine) [in same syringe]	IP	Once at induction, with additional doses as needed	Ketamine is a controlled substance . Xylazine is a controlled substance in Pennsylvania. Additional doses must be given at ½ strength.
Ketamine/Xylazine/Acepromazine	75-100 (ketamine) + 2-6 (xylazine) + 1-2 (acepromazine) [in same syringe]	IP	Once at induction, with additional doses as needed	Ketamine is a controlled substance . Xylazine is a controlled substance in Pennsylvania. Additional doses must be given at ½ strength.
Sodium Pentobarbital (Nembutal)	40-50	IP	Once at induction	Controlled substance . Recommended for terminal/acute procedures; must provide supplemental analgesic for invasive



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 7 of 14

				survival procedures. Difficult to obtain. Narrow margins of safety. Not recommended for survival surgery.
Propofol	10	IV	Once as induction before general anesthesia, or as a constant rate infusion	Respiratory depression is possible upon induction.
Urethane	1-2g/kg	IP	Once at induction	For non-survival procedures only. Non-pharmaceutical grade drug, therefore its use must be justified in the protocol due to scientific reasons. Due to potential human carcinogenicity, must be reviewed by the biosafety committee.

3.2 Anesthetic Reversal Agents

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Yohimbine	1.0-2.0	SC or IP	When needing to reverse the effects of xylazine	Yohimbine DOES NOT reverse the effects of Ketamine

3.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)

**Note: Prolonged use of NSAIDs may cause renal, gastrointestinal, or other problems*

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
-----------	--------------	-------	-----------	-------



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 8 of 14

Meloxicam (Metacam)	1	PO or SC	Once pre-operatively then every 12 to 24 hours as needed	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.
Carprofen (Rimadyl)	5	SC	Once pre-operatively then every 24 hours as needed	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.

3.4 Analgesics- Opioids (controlled substances)

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Buprenorphine	0.05	SC or IP	Once pre-operatively for preemptive analgesia, then every 6-8 hours	When used as sole analgesic, typical regimen is; once at time of procedure, second dose will be administered 4-6 hours later. Additional doses every 6-8 hrs as needed. Consider multi-modal analgesia with NSAID and local analgesic. Controlled substance
Buprenorphine Sustained-Release Lab (Bup SR Lab) 1mg/ml	0.5	SC	Once pre-operatively for preemptive analgesia	Provides 48 hours of pain relief; sedation may be seen for duration of effect. DO NOT DILUTE. Controlled substance.

3.5 Analgesics- Local Anesthetic/Analgesics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
-----------	--------------	-------	-----------	-------



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 9 of 14

Bupivacaine (Marcaine, Senosorcaine)	Dilute to 0.5%; do not exceed 7mg/kg total dose	SC or intradermal surrounding incision	Locally before making surgical incision, or before final skin closure	4-8 hour duration of action. Can be used with inhalants if other analgesics cannot be administered before animals awoken from isoflurane.
Lidocaine Hydrochloride	Dilute to 0.5%; do not exceed 7mg/kg total dose	SC or intradermal surrounding incision	Locally before making surgical incision, or before final skin closure	Faster onset than Bupivacaine, but short (<1 hour) duration of action. May be combined with bupivacaine.

Refer to ULAR Recommendations on Use of Bupivacaine for Perioperative Analgesia

[Bupivacaine \(Marcaine\) Recommendations for Use in Perioperative Analgesia](#)

4. Rabbits

4.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Isoflurane	4-5% (induction) 1-3% (maintenance)	Inhalant	Continuous	Rapid induction and recovery. Must provide supplemental analgesic for invasive survival procedures prior to beginning surgical procedure; must use precision vaporizer.
Ketamine/Xylazine	35-50 (ketamine) + 5-10 (xylazine) [in same syringe]	IM or SC	Once at induction, with additional doses as needed	Ketamine is a controlled substance . Xylazine is a controlled substance in Pennsylvania. IM injections of Ketamine often sting upon injection. Additional doses must be given at ½ strength.

4.2 Anesthetic Reversal Agents

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
-----------	--------------	-------	-----------	-------



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 10 of 14

Yohimbine	~0.2	IV or SC	When needing to reverse the effects of Xylazine	Yohimbine DOES NOT reverse the effects of Ketamine.
-----------	------	----------	---	---

4.3 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)

**Note: prolonged use of NSAIDs may cause renal, gastrointestinal, or other problems*

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Meloxicam (Metacam)	0.1-0.3	PO, IM, or SC	Once pre-operatively, then every 24 hours for 4 days	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.
Carprofen (Rimadyl)	4-5	SC	Once pre-operatively, then every 12-24 hours as needed	Depending on the procedure, may be used as a sole analgesic, or as multi-modal analgesia with buprenorphine.

4.4 Analgesics- Opioids (controlled substances)

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Buprenorphine	0.05-0.1	SC or IP	Once pre-operatively for preemptive analgesia and postoperatively every 4-12 hours	When used as sole analgesic, typical regimen is: once at time of procedure, second dose will be administered 4-6 hours later. Additional doses every 8-12 hours as needed. Consider multi-modal analgesia with NSAID and local analgesic.
Buprenorphine SR (3mg/ml)	0.1-0.3	SC	Once pre-operatively for preemptive analgesia	Provides 72 hours of pain relief. Controlled substance.



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 11 of 14

4.5 Analgesics- Opioids (controlled substances)

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Buprenorphine	0.05-0.1	SC	Once pre-operatively for preemptive analgesia and postoperatively every 4-12 hours	When used as sole analgesic, typical regimen is: once at time of procedure, second dose will be administered 4-6 hours later. Additional doses every 8-12 hours as needed. Consider multi-modal analgesia with NSAID and local analgesic. Controlled substance.
Fentanyl	25 mcg/hr for cats >3kg	Transdermal Patch	Once pre-operatively; change every 72 hours. Takes up to 12 hours to reach blood levels after patch placement	Can be given as sole analgesic 12 hours before surgery, or in conjunction with buprenorphine given at time of surgery. Controlled substance.

4.6 Analgesics- Local Anesthetic/Analgesics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Bupivacaine (Marcaine, Sensorcaine)	May dilute to 0.25%-0.5%; may be mixed in same syringe with lidocaine	SC or intradermal surrounding incision	Locally before making surgical incision, or before final skin closure	4-8 hour duration of action
Lidocaine Hydrochloride	May dilute to 0.5-1% (~10mg/ml); may be mixed in same syringe with bupivacaine	SC or intradermal surrounding incision	Locally before making surgical incision, or before skin closure	Faster onset than Bupivacaine, but short (<1 hour) duration of action

Refer to ULAR Recommendations on Use of Bupivacaine for Perioperative Analgesia

[Bupivacaine \(Marcaine\) Recommendations for Use in Perioperative Analgesia](#)



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 12 of 14

5. Swine

5.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Isoflurane	4-5% (induction)	Inhalant	Continuous	Rapid induction and recovery. Must provide supplemental analgesic for invasive survival procedures prior to beginning surgical procedure; must use precision vaporizer.
Telazol® Tiletamine	6-8	IM	Once at induction for sedation or preanesthetic	Telazol is a controlled substance and must be stored and refrigerated after reconstitution.
Propofol	16-22	IV	As induction agent, prior to general anesthesia or as Constant Rate Infusion	Respiratory depression upon induction is possible.

5.2 Analgesics- Non-Steroidal Anti-Inflammatories (NSAID)

**Note: prolonged use of NSAIDs may cause renal, gastrointestinal, or other problems*

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Meloxicam (Metacam)	0.1-0.3	PO, IM, or SC	Once pre-operatively then every 24 hours for up to 4 days	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.
Carprofen (Rimadyl)	4-5	SC	Once pre-operatively then every 12-24 hours as needed	Depending on the procedure, may be used as sole analgesic, or as multi-modal analgesia with buprenorphine.



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 13 of 14

5.3 Analgesics- Opioids (**controlled substances**)

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Buprenorphine	0.05-0.1	SC	Once pre-operatively for preemptive analgesia and postoperatively every 4-12 hours	When used as sole analgesic, typical regimen is: once at time of procedure, second dose will be administered 4-6 hours later. Additional doses every 8-12 hours as needed. Consider multi-modal analgesia with NSAID and local analgesic. Controlled substance.
Buprenorphine SR (3mg/ml)	0.1	SC	Once pre-operatively for preemptive analgesia	Provides 72 hours of pain relief. Controlled substance

6. Fish

6.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
Eugenol (Clove Oil)	40-100mg/L	Submersion	Continuous	Dosage depends on fish species and desired anesthetic depth.

7. Frogs

7.1 Anesthetics

Drug Name	Dose (mg/kg)	Route	Frequency	Notes
MS222	500-2000mg/L buffered with NaHCO ₃	Submersion	Continuous	Ensure pharmaceutical grade is used. Finquel®. Prepare in a fume hood as the powder is a respiratory irritant.



**Drug Formulary of Anesthetic and Analgesic Agents for
Laboratory Animals**

Document No.:	Edition No.:	Effective Date:	Page:
ACU-219	002	9/17/2024	Page 14 of 14

				https://www.drugs.com/vet/finquelms-222-tricaine-methanesulfonate.html
Benzocaine	200-300mg/L	Submersion	Continuous	

8. Resources

- Foley, Kendall, Turner. Clinical Management of Pain in Rodents. Comparative Medicine, Vol 69 (6) December 2019, p. 468-489.

9. Revisions

Edition 001/Effective Date: 11/11/2021 – Original Document

Edition 002/Revision Date: 9/11/2024 and Effective Date: 9/17/2024 – Revised Document.

- Updated formatting to new template.
- Listing xylazine as a controlled substance
- Removal of cat formulary